

Vertex covers in graphs with loops

MAURIZIO IMBESI

*Department of Mathematics, University of Messina
Viale F. Stagno d'Alcontres, 31 – I 98166 Messina, Italy
e-mail: imbesim@unime.it*

MONICA LA BARBIERA

*Department of Mathematics, University of Messina
Viale F. Stagno d'Alcontres, 31 – I 98166 Messina, Italy
e-mail: monicalb@unime.it*

Abstract

We investigate ideals of vertex covers for the edge ideals associated to considerable classes of connected graphs with loops and exhibit algebraic information about them, such as the existence of linear quotients, the computation of invariant values, and the Cohen-Macaulay property. These algebraic procedures are good instruments for evaluating situations of minimal node coverings in networks.

AMS 2010 Subject Classification: 05C25, 05C70, 13C14.

Key words and phrases: Graphs with loops. Linear quotients. Ideals of vertex covers. Cohen-Macaulay ideals.

Introduction

This article is devoted to study canonical properties of monomial ideals arising from remarkable classes of graphs with loops using computational and algebraic methods, in order to give analytic models of a network and of its connectivity. More precisely, the ideals of vertex covers for the edge ideals related to certain so-called graphs $\mathcal{K}'$ -type are examined and useful algebraic properties of them are considered.

Let $\mathcal{G}$ be any graph on vertex set $[n] = \{v_1, \dots, v_n\}$, set of edges $E(\mathcal{G})$, set of loops $L(\mathcal{G})$. An algebraic object attached to $\mathcal{G}$ is the edge ideal $I(\mathcal{G})$, a monomial ideal of the polynomial ring in n variables $R = K[X_1, \dots, X_n]$, K a field.

If $\mathcal{G}$ is a loopless graph, $I(\mathcal{G})$ is generated only by squarefree monomials $X_i X_j$

such that $\{v_i, v_j\} \in E(\mathcal{G})$, $i \neq j$. But, if $\mathcal{G}$ has loops, among the generators of $I(\mathcal{G})$ there are non-squarefree monomials X_i^2 such that $\{v_i, v_i\} \in L(\mathcal{G})$, for some $i = 1, \dots, n$.

The *ideal of vertex covers* $I_c(\mathcal{G})$ of $I(\mathcal{G})$ represents the algebraic transposition of the concept of (minimal) vertex cover of a graph $\mathcal{G}$. It is defined to be the ideal of R generated by all monomials $X_{i_1} \cdots X_{i_r}$ such that $(X_{i_1}, \dots, X_{i_r})$ is an associated minimal prime ideal of $I(\mathcal{G})$.

This terminology is intended to emphasize that a minimal prime "covers" or contains all the monomials in $I(\mathcal{G})$; sometimes, in particular situations, $I(\mathcal{G})$ reflects properties of $I_c(\mathcal{G})$, so that it is sufficient to study $I(\mathcal{G})$ for stating properties of $I_c(\mathcal{G})$.

In [7] it was introduced a significative class of edge ideals associated to connected loopless graphs $\mathcal{H}$ on vertex set $[n] = \{v_1, \dots, v_n\}$ formed by the union of a complete graph K_m , $m < n$, on vertex set $[m] = \{v_{\alpha_1}, \dots, v_{\alpha_m}\}$, $1 \leq \alpha_1 < \alpha_2 < \dots < \alpha_m = n$ and of star graphs with centers the vertices of K_m . Starting from $\mathcal{H}$ and adding loops on some nodes, it was also considered a larger class of connected graphs with loops $\mathcal{K}'$.

The ideals of vertex covers for the class of edge ideals associated to $\mathcal{H}$ and $\mathcal{K}'$ were described by using results about ideals of vertex covers for the edge ideals associated to the complete and the star graphs that make them up.

The explicit description of the generators of the ideals of vertex covers for the class of edge ideals associated to $\mathcal{K}'$ was obtained by generalizing a fundamental property about the structure of such ideals.

In the present paper we investigate the ideals $I_c(\mathcal{K}')$ in order to highlight some algebraic aspects of them.

In particular, we show in what cases the ideal of vertex covers $I_c(\mathcal{K}')$ admits linear quotients, and write them explicitly; we compute standard algebraic invariants of $I_c(\mathcal{K}')$ such as projective dimension, depth, Krull dimension, Castelnuovo-Mumford regularity; and we establish suitable conditions such that $I_c(\mathcal{K}')$ is a Cohen-Macaulay ideal. For details, see [3], [5], [6].

The study of such facts is devoted to find useful tools for improving actual critical situations for the connections in the field of communications and transport, specifically about minimal node covering.

We treat an optimization problem for the widespread deployment of police patrols on the main crossroads of a big city, in order to control all the streets and residential areas by standard routes.

We will show what should be the minimum number of patrols needed to cover adequately the areas around the intersections of the center and suburbs.

To perform this, a model of an appropriate graph with loops is given.

1 Ideals of vertex covers of graphs with loops

Let's recall algebraic definitions and notations.

Let $R = K[X_1, \dots, X_n]$ be a polynomial ring in n variables over an arbitrary field K such that $\deg X_i = 1$, for $i = 1, \dots, n$.

Let I be any monomial ideal of R .

$G(I)$ denotes the unique minimal set of monomial generators for I .

Definition 1.1. A *vertex cover* of $I \subset R$ is a subset C of $\{X_1, \dots, X_n\}$ such that each $u \in G(I)$ is divided by some $X_i \in C$.

C is called *minimal* if no proper subset of C is a vertex cover of I .

$h(I)$ denotes the minimal cardinality of the vertex covers of I .

Definition 1.2. $I \subset R$ is said to have *linear quotients* if there is an ordering $u_1, \dots, u_t$ of the monomials belonging to $G(I)$ such that the colon ideal $(u_1, \dots, u_{j-1}) : (u_j)$ is generated by a subset of $\{X_1, \dots, X_n\}$, for $j = 2, \dots, t$.

Remark 1.1. In [1], Conca and Herzog proved that an ideal generated in the same degree which has linear quotients admits a linear resolution.

Let $I \subset R$ have linear quotients with respect to the ordering $u_1, \dots, u_t$ of the monomials of $G(I)$. Let $q_j(I)$ be the number of the variables which is required to generate the ideal $(u_1, \dots, u_{j-1}) : (u_j)$.

Set $q(I) = \max_{2 \leq j \leq t} q_j(I)$.

Remark 1.2. The integer $q(I)$ is independent on the choice of the ordering of the generators that gives linear quotients ([4]).

Let's now give information on graphs and their ideals.

Let $\mathcal{G}$ be a connected graph on vertex set $[n] = \{v_1, \dots, v_n\}$. We indicate $E(\mathcal{G}) = \{\{v_i, v_j\} \mid v_i \neq v_j \text{ vertices of } \mathcal{G}\}$ the set of edges of $\mathcal{G}$ and $L(\mathcal{G}) = \{\{v_i, v_i\} \mid v_i \text{ any vertex of } \mathcal{G}\}$ the set of loops of $\mathcal{G}$. Hence $\{v_i, v_j\}$ is an edge joining v_i to v_j and $\{v_i, v_i\}$ is a loop of the vertex v_i .

If $L(\mathcal{G}) = \emptyset$, $\mathcal{G}$ is said a *simple* or *loopless* graph.

A *complete* graph on vertex set $[m] = \{v_1, \dots, v_m\}$, denoted by K_m , is a connected graph for which there exists an edge for all pairs $\{v_i, v_j\}$ of vertices of it. K'_m denotes a complete graph with loops on vertex set $[m]$.

A *star* graph on vertex set $[n] = \{v_i\} \cup \{v_1, \dots, v_{i-1}, v_{i+1}, v_n\}$ with center v_i , denoted by $\text{star}_i(n)$, $i = 1, \dots, n$, is a complete bipartite graph of the form $K_{1, n-1}$. $\text{star}'_i(n)$ denotes a star graph with loops on vertex set $[n]$ of center v_i .

If $R = K[X_1, \dots, X_n]$ such that each variable X_i corresponds to the vertex v_i of $\mathcal{G}$, the *edge ideal* $I(\mathcal{G})$ of $\mathcal{G}$ is the monomial ideal $(\{X_i X_j \mid \{v_i, v_j\} \in E(\mathcal{G})\} \cup \{X_i^2 \mid \{v_i, v_i\} \in L(\mathcal{G})\}) \subset R$.

Definition 1.3. A subset C of $[n] = \{v_1, \dots, v_n\}$ is said a *minimal vertex cover* of $\mathcal{G}$ if:

- (i) every edge of $\mathcal{G}$ is incident with one vertex in C , and
- (ii) no proper subset of C satisfies (i).

If C verifies condition (i) only, C is called a *vertex cover* of $\mathcal{G}$ and it is said to cover all the edges of $\mathcal{G}$.

The smallest number of vertices in any minimal vertex cover of $\mathcal{G}$ is said *vertex covering number*.

Remark 1.3. There exists a one to one correspondence between minimal vertex covers of $\mathcal{G}$ and minimal prime ideals of $I(\mathcal{G})$. In fact, $\wp$ is a minimal prime ideal of $I(\mathcal{G})$ if and only if $\wp = (C)$, for some minimal vertex cover C of $\mathcal{G}$. Thus $I(\mathcal{G})$ has primary decomposition $(C_1) \cap \dots \cap (C_r)$, where $C_1, \dots, C_r$ are the minimal vertex covers of $\mathcal{G}$.

Definition 1.4. The *ideal of vertex covers* of the edge ideal $I(\mathcal{G})$, denoted by $I_c(\mathcal{G})$, is the ideal of R generated by all the monomials $X_{i_1} \dots X_{i_r}$ such that $(X_{i_1}, \dots, X_{i_r})$ is an associated prime ideal of $I(\mathcal{G})$.

Note that if $\{v_i, v_i\} \in L(\mathcal{G})$ and v_i belongs to a vertex cover of the graph, $\{v_i, v_i\}$ can be thought to preserve the minimality.

The following generalizes the property for loopless graphs of [9]

Property 1.1. $I_c(\mathcal{G}) = \left(\bigcap_{\substack{\{v_i, v_j\} \in E(\mathcal{G}) \\ i \neq j}} (X_i, X_j) \right) \cap (X_k \mid \{v_k, v_k\} \in L(\mathcal{G}), k \neq i, j).$

From now on, we deal with the ideal of vertex covers for the class of connected graphs $\mathcal{K}'$ introduced in [7], namely the graphs $\mathcal{K}'$ -type.

More precisely, $\mathcal{K}'$ is the graph with loops on vertex set $[n] = \{v_1, \dots, v_n\}$ union of: (i) a complete graph $K_m, m < n$, with vertices $v_{\alpha_1}, \dots, v_{\alpha_m}$, $1 \leq \alpha_1 < \alpha_2 < \dots < \alpha_m = n$; (ii) star graphs $\text{star}_{\alpha_i}(\alpha_i - \alpha_{i-1})$ with vertices $v_{\alpha_{i-1}+1}, \dots, v_{\alpha_i}, \forall i = 1, \dots, m$, the index α_0 means 0.

The edge ideal of $\mathcal{K}'$ is:

$I(\mathcal{K}') = (X_{\alpha_1} X_{\alpha_2}, \dots, X_{\alpha_1} X_{\alpha_m}, X_{\alpha_2} X_{\alpha_3}, \dots, X_{\alpha_2} X_{\alpha_m}, \dots, X_{\alpha_{m-1}} X_{\alpha_m}, X_1 X_{\alpha_1}, X_2 X_{\alpha_1}, \dots, X_{\alpha_1-1} X_{\alpha_1}, X_{\alpha_1+1} X_{\alpha_2}, X_{\alpha_1+2} X_{\alpha_2}, \dots, X_{\alpha_2-1} X_{\alpha_2}, \dots, X_{\alpha_{m-1}+1} X_{\alpha_m}, \dots, X_{\alpha_m-1} X_{\alpha_m}, X_{t_1}^2, \dots, X_{t_l}^2) \subset R = K[X_1, \dots, X_n]$, with $\{t_1, \dots, t_l\} \subseteq \{1, \dots, n\}$.

Remark 1.4. $I(\mathcal{K}')$ has not linear quotients, because there is not an ordering of the monomials $f_1, \dots, f_s$ of $G(I(\mathcal{K}'))$ such that $(f_1, \dots, f_{j-1}) : (f_j)$ is generated by a subset of $\{X_1, \dots, X_n\}$, for $2 \leq j \leq s$. But there exists a subgraph $\mathcal{K}$ of $\mathcal{K}'$ whose loops lie only on the vertices belonging to K_m , such that $I(\mathcal{K})$ has linear quotients. According to Remark 1.1, $I(\mathcal{K})$ has a linear resolution. In general this is false for $I(\mathcal{K}')$.

The structure of the ideal of vertex covers for graphs $\mathcal{K}'$ -type was examined in [7], using the description of the ideals of vertex covers for the complete graph with loops and the star graph with loops that make it up.

Lemma 1.1. *Let K'_m be a complete graph on vertex set $[m] = \{v_1, \dots, v_m\}$ having loops on v_1, v_3, v_{m-1} . The ideal of vertex covers of $I(K'_m)$ is generated at most by $m - 1$ monomials. In particular,*

- (a) *if there are loops in all the vertices, $I_c(K'_m) = (X_1 X_2 \cdots X_m)$,*
- (b) *if there are loops in $r < m$ vertices, $v_{t_1}, \dots, v_{t_r}$, $\{t_1, \dots, t_r\} \subseteq \{1, \dots, m\}$, $I_c(K'_m)$ has $m - r$ generators and it is*
 $(\{X_{\sigma_1} \cdots X_{\sigma_{m-1}} \mid \sigma_j = t_j, \forall j = 1, \dots, r; \sigma_i \in \{1, \dots, m\} \setminus \{t_1, \dots, t_r\}, \forall i \neq j\})$.

Lemma 1.2. *Let $star'_n(n)$ be a star graph on vertex set $[n] = \{v_1, \dots, v_n\}$ having loops. The ideal of vertex covers of $I(star'_n(n))$ is generated at most by two monomials. In particular,*

- (a) *if the loops are in the vertices $v_1, \dots, v_{n-1}$, $I_c(star'_n(n)) = (X_1 \cdots X_{n-1})$,*
- (b) *if the loops are in v_3, v_{n-2} , $I_c(star'_n(n)) = (X_1 \cdots X_{n-1}, X_3 X_{n-2} X_n)$,*
- (c) *if there are loops in the center and in the vertices $v_{t_1}, \dots, v_{t_s}$, $\{t_1, \dots, t_s\} \subseteq \{1, \dots, n-1\}$, $I_c(star'_n(n)) = (X_{t_1} \cdots X_{t_s} X_n)$.*

The assertions of the preceding two lemmas descend from Property 1.1.

Theorem 1.1. *Let $\mathcal{K}'$ be as above and suppose that it has loops on some of its vertices, say $v_{\alpha_2}, v_{\alpha_4}, v_{\alpha_5}, v_{\alpha_{m-3}}, v_{\alpha_{m-1}}, v_{\alpha_{i-1}+j_1}, \dots, v_{\alpha_{i-1}+j_{p_i}}$, $i = 1, \dots, m$, where $\{j_1, \dots, j_{p_i}\} \subseteq \{1, \dots, \alpha_i - \alpha_{i-1} - 1\}$. The ideal of vertex covers $I_c(\mathcal{K}')$ has at most $m + 1$ monomial generators*

$$\begin{aligned} & X_{j_1} \cdots X_{j_{p_1}} X_{\alpha_1} X_{\alpha_1+j_1} \cdots X_{\alpha_1+j_{p_2}} X_{\alpha_2} X_{\alpha_2+j_1} \cdots X_{\alpha_{m-1}+j_{p_m}} X_{\alpha_m}, \\ & X_1 \cdots X_{\alpha_1-1} X_{\alpha_1+j_1} \cdots X_{\alpha_1+j_{p_2}} X_{\alpha_2} X_{\alpha_2+j_1} \cdots X_{\alpha_{m-1}+j_{p_m}} X_{\alpha_m}, \\ & X_{j_1} \cdots X_{j_{p_1}} X_{\alpha_1} X_{\alpha_1+1} \cdots X_{\alpha_2-1} X_{\alpha_2} X_{\alpha_2+j_1} \cdots X_{\alpha_{m-1}+j_{p_m}} X_{\alpha_m}, \\ & \dots, X_{j_1} \cdots X_{\alpha_{m-3}+j_{p_{m-2}}} X_{\alpha_{m-2}} X_{\alpha_{m-2}+1} \cdots \\ & X_{\alpha_{m-1}-1} X_{\alpha_{m-1}} X_{\alpha_{m-1}+j_1} \cdots X_{\alpha_{m-1}+j_{p_m}} X_{\alpha_m}, \\ & X_{j_1} \cdots X_{\alpha_{m-2}+j_{p_{m-1}}} X_{\alpha_{m-1}} X_{\alpha_{m-1}+1} \cdots X_{\alpha_m-1}. \end{aligned}$$

Significative particular cases for determining the structure of the ideal of vertex covers related to a graph $\mathcal{K}'$ -type are (see [7]):

- (a) some vertices of K_m are not centers of star graphs,
- (b) the loops lie only on the vertices of K_m , and
- (c) the loops lie only on the vertices not belonging to K_m .

Example 1.1. Let $\mathcal{K}'$ be the connected graph on vertex set $\{v_1, \dots, v_{11}\}$ given by $K_3 \cup \text{star}_4(4) \cup \text{star}_8(4) \cup \text{star}_{11}(3) \cup \{v_2, v_2\} \cup \{v_5, v_5\} \cup \{v_7, v_7\} \cup \{v_{11}, v_{11}\}$. The ideal of vertex covers for $\mathcal{K}'$ is $I_c(\mathcal{K}') = (X_2X_4X_5X_6X_7X_{11}, X_2X_4X_5X_7X_8X_{11}, X_1X_2X_3X_5X_7X_8X_{11})$.

2 Linear quotients and properties for $I_c(\mathcal{K})'$

In this section we illustrate some algebraic aspects of $I_c(\mathcal{K}')$.

We prove that there exists an ordering on the number of variables of every monomial of $I_c(\mathcal{K}')$ for which this ideal admits linear quotients and write them explicitly; we also compute standard algebraic invariants of $I_c(\mathcal{K}')$ such as projective dimension, depth, Krull dimension, Castelnuovo-Mumford regularity; and finally, because the ideals of vertex covers $I_c(\mathcal{H})$ are Cohen-Macaulay ([8]), we show that this property is preserved for the ideals of vertex covers $I_c(\mathcal{K}')$ provided that the graphs $\mathcal{K}'$ -type have loops on all the vertices corresponding at least to a monomial generator of $I_c(\mathcal{H})$.

Let's examine the existence of linear quotients for the ideal of vertex covers $I_c(\mathcal{K}')$.

Theorem 2.1. *Let $\mathcal{K}'$ be as above, but at most $m-2$ vertices of K_m have loops. Then there exists an order $\leq$ on the number of variables in any monomial of $I_c(\mathcal{K}')$, natural order of indices for the same number of variables, such that $I_c(\mathcal{K}')$ has linear quotients.*

Proof. Theorem 1.1 and its consequences give us the minimal set of monomial generators for the ideal of vertex covers $I_c(\mathcal{K}')$. Such generators are squarefree and may have different degrees. Put them in increasing order according to the degrees and, for the same degree, the leftmost nonzero difference of the indices be negative.

We have to distinguish two cases and for each of them will explicitly write the linear quotients of $I_c(\mathcal{K}')$.

Case I. — The vertices $v_{\alpha_1}, \dots, v_{\alpha_m}$ of K_m in $\mathcal{K}'$ are centers of star graphs with or without loops, otherwise there is a loop on any vertex of K_m that is not center of a star graph.

▷ There are no loops on the vertices of K_m . The linear quotients are:

- $(X_{\alpha_1}), \dots, (X_{\alpha_m})$
if there are ends not covered by loops for each star graph;

- $(X_{\alpha_1}), \dots, (X_{\alpha_{m-1}})$
if star graph with center in v_{α_m} has loops in all its remaining vertices.
- ▷ There is a loop on the vertex v_{α_m} of K_m . The linear quotients are:
 - $(X_{\alpha_1}), \dots, (X_{\alpha_{m-1}})$
if there are ends not covered by loops for each star graph whose center has not a loop, or v_{α_m} is not center of a star graph;
 - $(X_{\alpha_1}), \dots, (X_{\alpha_{m-2}})$
if star graph with center in $v_{\alpha_{m-1}}$ has loops in all its remaining vertices.
- ▷ There are loops on vertices $v_{\alpha_{m-1}}, v_{\alpha_m}$ of K_m . The linear quotients are:
 - $(X_{\alpha_1}), \dots, (X_{\alpha_{m-2}})$
if there are ends not covered by loops for each star graph whose center has not a loop, or $v_{\alpha_{m-1}}, v_{\alpha_m}$ are not centers of star graphs;
 - $(X_{\alpha_1}), \dots, (X_{\alpha_{m-3}})$
if star graph with center in $v_{\alpha_{m-2}}$ has loops in all its remaining vertices.

.....
- ▷ There are loops on vertices $v_{\alpha_3}, \dots, v_{\alpha_m}$ of K_m . The linear quotients are:
 - $(X_{\alpha_1}), (X_{\alpha_2})$
if there are ends not covered by loops for each star graph whose center has not a loop, or $v_{\alpha_3}, \dots, v_{\alpha_m}$ are not centers of star graphs;
 - (X_{α_1})
if star graph with center in v_{α_2} has loops in all its remaining vertices.

Observe that if there are loops on $m-1$ or m vertices of K_m , $I_c(\mathcal{K}')$ has not linear quotients because it is generated by a unique monomial whose variables correspond to the vertices with loops of $\mathcal{K}'$.

Case II. — The vertices without loops of K_m in $\mathcal{K}'$ are not all centers of star graphs with or without loops.

- ▷ There are no loops on the vertices of K_m . The linear quotients are:
 - $(X_{\alpha_{i_1}}), \dots, (X_{\alpha_{i_{m-1}}})$
 $\forall i_1 \neq \dots \neq i_{m-1} \in \{\alpha_1, \alpha_2, \dots, \alpha_m\}$.
- ▷ There is a loop on the vertex v_{α_m} that can be center of a star graph with or without loops. The linear quotients are:

- $(X_{\alpha_{i_1}}, \dots, (X_{\alpha_{i_{m-2}}})$
when none, some or all the vertices $v_{\alpha_{i_1}}, \dots, v_{\alpha_{i_{m-2}}}$ of K_m are centers of star graphs with or without loops, $\forall i_1 \neq \dots \neq i_{m-2} \in \{\alpha_1, \dots, \alpha_{m-1}\}$.
- ▷ There are loops on vertices $v_{\alpha_{m-1}}, v_{\alpha_m}$ and each one can be center of a star graph with or without loops. The linear quotients are:
 - $(X_{\alpha_{i_1}}, \dots, (X_{\alpha_{i_{m-3}}})$
when none, some or all the vertices $v_{\alpha_{i_1}}, \dots, v_{\alpha_{i_{m-3}}}$ of K_m are centers of star graphs with or without loops, $\forall i_1 \neq \dots \neq i_{m-3} \in \{\alpha_1, \dots, \alpha_{m-2}\}$.
 -
- ▷ There are loops on vertices $v_{\alpha_4}, \dots, v_{\alpha_m}$ and each one can be center of a star graph with or without loops. The linear quotients are:
 - $(X_{\alpha_{i_1}}, (X_{\alpha_{i_2}})$
when none, one or both the vertices $v_{\alpha_{i_1}}, v_{\alpha_{i_2}}$ of K_m are centers of star graphs with or without loops, $\forall i_1 \neq i_2 \in \{\alpha_1, \alpha_2, \alpha_3\}$.
- ▷ There are loops on vertices $v_{\alpha_3}, \dots, v_{\alpha_m}$ and each one can be center of a star graph with or without loops. The linear quotients are:
 - (X_{α_i}) when $v_{\alpha_i} \in K_m$ is or not center of a star graph with or without loops, $\forall i \in \{\alpha_1, \alpha_2\}$.

As previously observed, if there are loops on at least $m-1$ vertices of K_m , $I_c(\mathcal{K}')$ has not linear quotients. $\square$

Remark 2.1. If $\mathcal{K}'$ satisfies the hypotheses of Theorem 2.1, $I_c(\mathcal{K}')$ has linear quotients. By Theorem 1.1, $I_c(\mathcal{K}')$ is not necessarily generated in the same degree. According to Remark 1.1, $I_c(\mathcal{K}')$ has not in general a linear resolution.

The existence of linear quotients for a monomial ideal $I \subset R$ makes easier the computation of some algebraic invariants of R/I . In fact :

- ◊ the *projective dimension*, $\text{pd}_R(R/I) = q(I) + 1$,
- ◊ the *depth*, $\text{depth}_R(R/I) = \dim R - \text{pd}_R(R/I)$,
where $\dim R = \dim_R(R/I) + h(I)$ is the *Krull dimension*,
- ◊ the *Castelnuovo-Mumford regularity*,
 $\text{reg}_R(R/I) = \max\{\deg f \mid f \text{ minimal generator of } I\} - 1$.

Lemma 2.1. Let $R = K[X_1, \dots, X_n]$ and $I_c(\mathcal{K}') \subset R$. Then $h(I_c(\mathcal{K}')) = 1$.

Proof. Being $I_c(\mathcal{K}')$ an ideal of vertex covers associated to a graph with at least a loop v_i , the variable X_i is common in all its generators. So there exists a vertex cover of $I_c(\mathcal{K}')$ having the unique element X_i , namely a set of minimal cardinality among the vertex covers of $I(\mathcal{K}')$. $\square$

Lemma 2.2. *Let $R = K[X_1, \dots, X_n]$ and $I_c(\mathcal{K}') \subset R$. Suppose that $I_c(\mathcal{K}')$ has linear quotients. Then $q(I_c(\mathcal{K}')) = 1$.*

Proof. As a consequence of Theorem 2.1, $I_c(\mathcal{K}')$ has linear quotients if it is generated at least by two monomials. In this case, the number of variables for generating the ideal $(f_1, \dots, f_{h-1}) : (f_h)$ is 1, for all $h = 1, \dots, t$, $t \leq m+1$, m the number of vertices in K_m . $\square$

Let's study algebraic invariants values for the ideal of vertex covers $I_c(\mathcal{K}')$.

Theorem 2.2. *Let $\mathcal{K}'$ be as above. Let σ denote the maximal number of vertices among the star graphs of $\mathcal{K}'$. Let $R = K[X_1, \dots, X_n]$ and $I_c(\mathcal{K}') \subset R$ be the ideal of vertex covers related to $\mathcal{K}'$. If $I_c(\mathcal{K}')$ has linear quotients, then:*

- 1) $\text{pd}_R(R/I_c(\mathcal{K}')) = 2$.
- 2) $\text{depth}_R(R/I_c(\mathcal{K}')) = n - 2$.
- 3) $\dim_R(R/I_c(\mathcal{K}')) = n - 1$.
- 4) $(m-1) + (\sigma-2) \leq \text{reg}_R(R/I_c(\mathcal{K}')) \leq n-2$.

Proof. 1) The length of a minimal free resolution of $R/I_c(\mathcal{K}')$ over R is equal to $q(I_c(\mathcal{K}')) + 1$ ([4], Corollary 1.6). So, by Lemma 2.2, $\text{pd}_R(R/I_c(\mathcal{K}')) = 2$.
2) As a consequence of 1), by Auslander-Buchsbaum formula, one has that $\text{depth}_R(R/I_c(\mathcal{K}')) = n - \text{pd}_R(R/I_c(\mathcal{K}')) = n - 2$.
3) It results $\dim_R(R/I_c(\mathcal{K}')) = \dim_R R - h(I_c(\mathcal{K}'))$ ([2]). Hence, by Lemma 2.1, $\dim_R(R/I_c(\mathcal{K}')) = n - 1$.
4) By [1], Lemma 4.1, $\text{reg}(I_c(\mathcal{K}'))$ is the maximum of the degrees for any minimal generator of $I_c(\mathcal{K}')$. By hypothesis, $I_c(\mathcal{K}')$ cannot be generated by a unique monomial, so any element of it has at most $n-1$ variables. Indeed, the lower bound is given by the maximal degree of the generators of the correspondent $I_c(\mathcal{H})$ ([7], Prop. 2.1, Thm. 2.1). $\square$

If $I_c(\mathcal{K}')$ is generated by a unique monomial, it has not linear quotients. However the following result holds:

Corollary 2.1. *Let $R = K[X_1, \dots, X_n]$ and $I_c(\mathcal{K}') \subset R$. If $I_c(\mathcal{K}')$ is generated by a unique monomial, then:*

- 1) $\text{pd}_R(R/I_c(\mathcal{K}')) = 1$.
- 2) $\text{depth}_R(R/I_c(\mathcal{K}')) = n - 1$.
- 3) $\dim_R(R/I_c(\mathcal{K}')) = n - 1$.
- 4) $\text{reg}_R(R/I_c(\mathcal{K}')) \leq n - 1$.

Example 2.1. Let $\mathcal{K}' = K_5 \cup \text{star}_3(3) \cup \text{star}_6(3) \cup \text{star}_8(2) \cup \text{star}_{12}(3) \cup \{v_5, v_5\} \cup \{v_8, v_8\} \cup \{v_{12}, v_{12}\}$ on vertex set $\{v_1, \dots, v_{12}\}$.
 $I_c(\mathcal{K}') = (X_3X_5X_6X_8X_{12}, X_3X_4X_5X_8X_9X_{12}, X_1X_2X_5X_6X_8X_9X_{12})$.
 $I_c(\mathcal{K}')$ has linear quotients $(X_3), (X_6)$, but it is not generated in the same degree so it has not a linear resolution. In fact :

$$0 \rightarrow R(-7) \oplus R(-8) \rightarrow R(-5) \oplus R(-6) \oplus R(-7) \rightarrow R/I_c(\mathcal{K}') \rightarrow R \rightarrow 0.$$

A computation for getting algebraic invariants gives :

- 1) $\text{pd}_R(R/I_c(\mathcal{K}')) = 2$.
- 2) $\text{depth}_R(R/I_c(\mathcal{K}')) = 10$.
- 3) $\dim_R(R/I_c(\mathcal{K}')) = 11$.
- 4) $\text{reg}_R(R/I_c(\mathcal{H})) = 6$.

About the Cohen-Macaulay property for $I_c(\mathcal{K}')$.

Recall that an ideal $I \subset R$ is *Cohen-Macaulay* if and only if R/I is Cohen-Macaulay, i.e $\text{depth}(R/I) = \dim(R/I)$.

Starting from some characteristics of the loopless graph $\mathcal{H}$, formed by the union of a complete graph K_m , $m < n$, together with star graphs with centers the vertices of K_m , and from the structure of the edge ideal $I(\mathcal{H})$ of $\mathcal{H}$, for the ideal of vertex covers $I_c(\mathcal{H})$ the following holds.

Proposition 2.1. $I_c(\mathcal{H})$ is a Cohen-Macaulay ideal.

Proof. See [8], Proposition 2.4. □

Next result shows that such a property is preserved for the ideals $I_c(\mathcal{K}')$ related to the graphs $\mathcal{K}'$ -type which have loops on all the vertices corresponding at least to a monomial generator of $I_c(\mathcal{H})$.

Theorem 2.3. Let $I_c(\mathcal{H})$ be the ideal of vertex covers for the edge ideal associated to the graph $\mathcal{H}$ and $f_1, \dots, f_t$, $t \leq m-1$ be its monomial generators. Let's consider the graph $\mathcal{K}'$ obtained from $\mathcal{H}$ by inserting loops at least in the vertices that correspond to all the variables on any generator f_i of $I_c(\mathcal{H})$. Then the ideal of vertex covers $I_c(\mathcal{K}')$ is Cohen-Macaulay.

Proof. In [7] the structure of the generators of the ideal of vertex covers $I_c(\mathcal{H})$ was studied. When a loop $\{v_i, v_i\}$ is added to the graph $\mathcal{H}$, all the generators of the ideal of vertex covers for the edge ideal associated to the new graph $\mathcal{K}'$ will have the common variable X_i . If the loops added are relative at least to all the variables in any generator of $I_c(\mathcal{H})$, the ideal of vertex covers $I_c(\mathcal{K}')$ will be generated by a unique monomial. The assertion follows from Corollary 2.1, because $\text{depth}_R(R/I_c(\mathcal{K}')) = \dim_R(R/I_c(\mathcal{K}'))$. □

Example 2.2. Let $\mathcal{H} = K_5 \cup \text{star}_3(3) \cup \text{star}_6(3) \cup \text{star}_8(2) \cup \text{star}_{12}(3)$ on vertex set $\{v_1, \dots, v_{12}\}$. $I_c(\mathcal{H}) = (X_1X_2X_6X_8X_9X_{12}, X_3X_4X_5X_8X_9X_{12}, X_3X_6X_7X_9X_{12}, X_3X_6X_8X_9X_{10}X_{11}, X_3X_6X_8X_{12})$.

Let $\mathcal{K}' = K_5 \cup \text{star}_3(3) \cup \text{star}_6(3) \cup \text{star}_8(2) \cup \text{star}_{12}(3) \cup \{v_2, v_2\} \cup \{v_3, v_3\} \cup \{v_4, v_4\} \cup \{v_5, v_5\} \cup \{v_8, v_8\} \cup \{v_9, v_9\} \cup \{v_{10}, v_{10}\} \cup \{v_{12}, v_{12}\}$ be the graph obtained from $\mathcal{H}$ by inserting loops on the vertices correspondent to the monomial generator $X_3X_4X_5X_8X_9X_{12}$ of $I_c(\mathcal{H})$ and in other ones.

It is $I_c(\mathcal{K}') = (X_2X_3X_4X_5X_8X_9X_{10}X_{12})$.

One has: $\text{depth}_R(R/I_c(\mathcal{K}')) = 11 = \dim_R(R/I_c(\mathcal{K}'))$.

Then $I_c(\mathcal{K}')$ is Cohen-Macaulay.

Last theorem extends to any connected graph with loops, as the following formulates

Theorem 2.4. *Let $\mathcal{G}$ be a connected loopless graph on vertex set $[n] = \{v_1, \dots, v_n\}$ and edge ideal $I(\mathcal{G}) \subset R = K[X_1, \dots, X_n]$. Let $I_c(\mathcal{G})$ be the ideal of vertex covers for the edge ideal associated to $\mathcal{G}$ and suppose that $I_c(\mathcal{G})$ is Cohen-Macaulay. Let's consider the graph $\mathcal{G}'$ built from $\mathcal{G}$ by adding loops at least on the vertices that correspond to all the variables in any generator of $I_c(\mathcal{G})$. Then the ideal of vertex covers $I_c(\mathcal{G}')$ is Cohen-Macaulay.*

Proof. A generalization of the proof of Theorem 2.3, bearing in mind that the hypotheses make $I_c(\mathcal{G}')$ a principal ideal. $\square$

3 An optimization problem of police control

Let's evaluate an application for the widespread deployment of a minimum number of police patrols in crucial crossroads or squares of a big city, in order to control all the streets and residential areas by standard routes.

To perform this, we can think for modeling the city through a graph $\mathcal{K}'$ -type. To be more precise :


- vertices represent crucial nodes, such as crossroads or squares ;
- every edge can be seen as the route assigned to the patrol stationed in a vertex to reach an adjacent one ;
- every loop can be intended as the route assigned to the patrol to return to the starting vertex without going through anyone else.

The least number of patrols needed to adequately cover the areas around the intersections of the center and suburbs is given by the monomials with the minimum number of variables which generate the ideal of vertex covers of the edge ideal related to $\mathcal{K}'$.

They just identify the crossroads or squares that have to be monitored. If there are different generators of the ideal of vertex covers with the smallest number of variables, there will be several solutions to the minimum problem.

Remark 3.1. If the ideal of vertex covers for $\mathcal{K}'$ has only one generator, then it is Cohen-Macaulay. This can be interpreted thinking of the graphs $\mathcal{K}'$ -type have a unique covering, namely one solution to the problem of minimal vertex covers.

Example 3.1. Let's suppose the map of any city can be modeled through the graph $\mathcal{K}'$ -type below



The minimum number of police patrols necessary for covering the main crossroads or squares, represented by the vertices of the graph, in order to control all the streets and residential areas by standard routes, outlined by edges and loops of it, is linked to the generators with the minimum number of variables of the ideal of vertex covers $I_c(\mathcal{K}')$.

According to Theorem 1.1, a computation yields the displayed monomial generators for it:

$$\begin{aligned} &X_1 X_2 X_3 X_7 X_9 X_{10} X_{12} X_{14} X_{15} X_{16} X_{17} X_{19} X_{20} X_{23} X_{25} X_{26} X_{27} X_{28} X_{29} X_{30} X_{32} X_{33}, \\ &X_2 X_3 X_4 X_7 X_9 X_{10} X_{11} X_{12} X_{13} X_{15} X_{16} X_{17} X_{19} X_{20} X_{23} X_{25} X_{26} X_{27} X_{28} X_{29} X_{30} X_{32} X_{33}, \\ &X_2 X_3 X_4 X_7 X_9 X_{10} X_{12} X_{14} X_{15} X_{16} X_{17} X_{18} X_{20} X_{21} X_{22} X_{25} X_{26} X_{27} X_{28} X_{29} X_{30} X_{32} X_{33}, \\ &X_2 X_3 X_4 X_7 X_9 X_{10} X_{12} X_{14} X_{15} X_{16} X_{17} X_{18} X_{20} X_{23} X_{25} X_{26} X_{27} X_{28} X_{29} X_{30} X_{32} X_{33}, \\ &X_2 X_3 X_4 X_7 X_9 X_{10} X_{12} X_{14} X_{15} X_{16} X_{17} X_{19} X_{20} X_{21} X_{22} X_{25} X_{26} X_{27} X_{28} X_{29} X_{30} X_{32} X_{33}, \\ &X_2 X_3 X_4 X_7 X_9 X_{10} X_{12} X_{14} X_{15} X_{16} X_{17} X_{19} X_{20} X_{23} X_{25} X_{26} X_{27} X_{28} X_{29} X_{32} X_{33}. \end{aligned}$$

They correspond to minimal vertex covers of $\mathcal{K}'$. The only one with the

smallest number of vertices is:

$$C = \{2, 3, 4, 7, 9, 10, 12, 14, 15, 16, 17, 19, 20, 23, 25, 26, 27, 28, 29, 32, 33\}.$$

Thus, the requested minimum number of patrols is the vertex covering number of $\mathcal{K}'$, namely twenty-one; they have to be set out on the nodes of C .

References

- [1] Conca, A., and Herzog, J. - Castelnuovo-Mumford regularity of product of ideals. *Collect. Math.* **54** (2003), 137–152.
- [2] Herzog, J., and Hibi, T. - Cohen-Macaulay polymatroidal ideals. *Eur. J. Comb.* **27**, 4 (2006), 513–517.
- [3] Herzog, J., and Hibi, T. - Monomial Ideals. *Springer, Graduate Texts in Mathematics* **260** (2011).
- [4] Herzog, J., and Takayama, Y. - Resolutions by mapping cones. *Homology, Homotopy Appl.* **4**, 2 (2002), 277–294.
- [5] Imbesi, M., and La Barbiera M. - Invariants of symmetric algebras associated to graphs. *Turkish J. Math.*, (2011) In press.
- [6] Imbesi, M., and La Barbiera M. - Algebraic properties of non-squarefree graph ideals. *Math. Reports*, (2011), In press.
- [7] Imbesi, M., and La Barbiera M. - Monomial ideals of graphs with loops. *Anal. Stiinț. Univ. Al. I. Cuza din Iași, Sect. Mat.*, (2012), In press
- [8] Imbesi, M., and La Barbiera M. - Vertex covers and sensor network. (2012), Submitted.
- [9] Villarreal, R.H. - Monomial Algebras. *M. Dekker Inc., New York, Pure and Appl. Math.* **238** (2001).